

# Derivation Tree Analysis for Accelerated Fixed-Point Computation

Javier Esparza, Stefan Kiefer, Michael Luttenberger

Institut für Informatik, Technische Universität München, 85748 Garching, Germany  
{esparza, kiefer, luttenbe}@model.in.tum.de

**Abstract.** We show that for several classes of idempotent semirings the least fixed-point of a polynomial system of equations  $X = f(X)$  is equal to the least fixed-point of a *linear* system obtained by “linearizing” the polynomials of  $f$  in a certain way. Our proofs rely on derivation tree analysis, a proof principle that combines methods from algebra, calculus, and formal language theory, and was first used in [5] to show that Newton’s method over commutative and idempotent semirings converges in a linear number of steps. Our results lead to efficient generic algorithms for computing the least fixed-point. We use these algorithms to derive several consequences, including an  $O(N^3)$  algorithm for computing the throughput of a context-free grammar (obtained by speeding up the  $O(N^4)$  algorithm of [2]), and a generalization of Courcelle’s result stating that the downward-closed image of a context-free language is regular [3].

## 1 Introduction

Systems  $X = f(X)$  of fixed-point equations, where  $f$  is a system of polynomials, appear naturally in semantics, interprocedural program analysis, language theory, and in the study of probabilistic systems (see e.g. [6, 7, 9, 12]). In all these applications the equations are interpreted over  $\omega$ -continuous semirings, an algebraic structure that guarantees the existence of a least solution  $\mu f$ . The key algorithmic problem is to compute or at least approximate  $\mu f$ .

In [5, 4] we generalized Newton’s method—the well-known method of numerical mathematics for approximating a zero of a differentiable function—to arbitrary  $\omega$ -continuous semirings. Given a polynomial system  $f$ , our generalized method computes a sequence of increasingly accurate approximations to  $\mu f$ , called Newton approximants. We showed in [5] that the  $n$ -th Newton approximant of a system of  $n$  equations over an idempotent (w.r.t. addition) and commutative (w.r.t. multiplication) semiring is already equal to  $\mu f$ . This theorem leads to a generic computing procedure.

Our proof of this result uses a (to the best of our knowledge) novel technique, which we call *derivation tree analysis*. The system  $f$  induces a set  $\mathcal{T}$  of *derivation trees*, a generalization of the well-known derivation trees of context-free grammars. Each tree can be naturally assigned a semiring element, called the *yield* of the tree. It is easy to show that  $\mu f$  is equal to the sum of the yields of all derivation trees. Derivation tree analysis first identifies a subset  $T'$  of derivation trees whose total yield  $Y(T')$  is easy to compute in some sense, and then proves that  $T'$  satisfies the *embedding property*:  $Y(t) \sqsubseteq Y(T')$  for every derivation tree  $t$ . If the semiring is idempotent, the embedding

property implies  $Y(T) = Y(T')$ , and so  $\mu f = Y(T')$ . In [5], the set  $T'$  was chosen so that  $Y(T')$  is equal to the  $n$ -th Newton approximant, and the embedding property was proved using some tree surgery and exploiting the commutativity of the semiring.

The computation of the  $n$ -th Newton approximant can still require considerable resources. In this paper we present a further application of derivation tree analysis to idempotent semirings, leading to more efficient computation algorithms. For this, we define the set  $\mathcal{B}$  of *bamboos* of a system  $f$ . Loosely speaking, bamboos are derivation trees with an arbitrarily long stem but only short branches. We first show that  $Y(\mathcal{B})$  is the solution of a linear system of equations whose functions are similar (but not identical) to the straightforward linearisation of  $f$ . Then, we prove that the following three classes of semirings satisfy the embedding property:

- *Star-distributive semirings* are idempotent and commutative semirings satisfying the additional axiom  $(a+b)^* = a^* + b^*$  (where  $*$  is the well-known Kleene iteration operator). The so-called “tropical”  $(\min, +)$ -semiring over the reals (extended with  $+\infty$  and  $-\infty$ ) is star-distributive. Our tree analysis leads to an algorithm for computing  $\mu f$  very similar to the generalized Bellman-Ford algorithm of Gawlitza and Seidl [8]. We use it to derive a new algorithm for computing the throughput of a context-free grammar, a problem introduced and analyzed by Caucal et al. in [2]. Our algorithm runs in  $O(N^3)$ , a factor  $N$  faster than the algorithm presented in [2].

- *Lossy semirings* are idempotent semirings satisfying the additional axiom  $a + 1 = a$  where  $1$  is the neutral element of multiplication. A natural model are downward-closed languages with union and concatenation as operations. Lossy semirings find application in the verification of lossy channel systems, a model of computation thoroughly investigated by Abdulla et al. (see e.g. [1]). Our tree analysis leads to an algebraic proof of Courcelle’s theorem stating that the downward closure of a context-free language is effectively regular [3].

- *1-bounded semirings* are idempotent semirings where the equation  $a + 1 = 1$  holds. A natural example is the “maximum probability” semiring with the interval  $[0, 1]$  as carrier, maximum as addition, and standard multiplication over the reals. Using derivation tree analysis it is very easy to show that the least fixed-point  $\mu f$  of a polynomial system  $f$  with  $n$  variables is given by  $f^n(\mathbf{0})$ , the  $n$ -fold application of  $f$  to  $\mathbf{0}$ .

The rest of the paper is organized as follows. After the preliminaries in Section 2 we introduce derivation tree analysis in Section 3. Bamboos are defined in Section 4. In the Sections 5, 6 and 7 we apply derivation tree analysis to the semiring classes mentioned above. An appendix includes the missing proofs.

## 2 Preliminaries

As usual,  $\mathbb{N}$  denotes the set of natural numbers including 0.

An *idempotent semiring*  $\mathcal{S} = \langle S, +, \cdot, 0, 1 \rangle$  consists of a commutative, idempotent additive monoid  $\langle S, +, 0 \rangle$ , and a multiplicative monoid  $\langle S, \cdot, 1 \rangle$ . In the following we often omit the dot  $\cdot$  in products. Both algebraic structures are connected by left- and right-distributivity, e.g.  $a(b + c) = ab + ac$ , and by the requirement that  $0 \cdot a = 0$  for all  $a \in S$ . The *natural-order relation*  $\sqsubseteq S \times S$  is defined by  $a \sqsubseteq b \Leftrightarrow a + b = b$ . The semiring  $\mathcal{S}$  is *naturally ordered* if  $\sqsubseteq$  is a partial order.

An idempotent, naturally ordered semiring  $\mathcal{S}$  is  $\omega$ -continuous, if countable summation  $\sum_{i \in \mathbb{N}} a_i \in \mathcal{S}$  is defined (with  $a_i \in \mathcal{S}$ ), and satisfies the following requirements: (i) summation is continuous, i.e.,  $\sup^{\sqsubseteq} \{a_0 + a_1 + \dots + a_k \mid k \in \mathbb{N}\} = \sum_{i \in \mathbb{N}} a_i$  for all sequences  $a : \mathbb{N} \rightarrow \mathcal{S}$ ; (ii) distributivity extends in the natural way to countable summation; and (iii)  $\sum_{j \in J} \sum_{i \in I_j} a_i = \sum_{i \in \mathbb{N}} a_i$  holds for all partitions  $(I_j)_{j \in J}$  of  $\mathbb{N}$ . In every such  $\omega$ -continuous semiring the Kleene-star operator  $*$  :  $\mathcal{S} \rightarrow \mathcal{S}$  is well-defined by  $a^* := \sum_{k \in \mathbb{N}} a^k$  for all  $a \in \mathcal{S}$ . In the following we consider only idempotent  $\omega$ -continuous semirings  $\mathcal{S}$ . We refer to them as *io-semirings*.

We fix a finite, non-empty set  $\mathcal{X}$  of *variables* for the rest of the section, and use  $n$  to denote  $|\mathcal{X}|$  in the following. A map from  $\mathcal{X}$  to  $\mathcal{S}$  is called a *vector*. The set of all vectors is denoted by  $V$ . We write both  $v(X)$  and  $v_X$  for the value of a vector  $v$  at  $X \in \mathcal{X}$ , also called the  $X$ -component of  $v$ . Sum of vectors is defined componentwise: given a countable set  $I$  and a vector  $v_i$  for every  $i \in I$ , we denote by  $\sum_{i \in I} v_i$  the vector given by  $(\sum_{i \in I} v_i)(X) = \sum_{i \in I} v_i(X)$  for every  $X \in \mathcal{X}$ .

A *monomial of degree  $k$*  is a finite expression  $a_1 X_1 a_2 \dots a_k X_k a_{k+1}$  where  $k \geq 0$ ,  $a_1, \dots, a_{k+1} \in \mathcal{S} \setminus \{0\}$  and  $X_1, \dots, X_k \in \mathcal{X}$ . A *polynomial* is an expression of the form  $m_1 + \dots + m_k$  where  $k \geq 0$  and  $m_1, \dots, m_k$  are monomials. Since  $\mathcal{S}$  is idempotent, we assume w.l.o.g. that all monomials of a polynomial are distinct. The degree of a polynomial is the largest degree of its monomials. We let  $\mathcal{S}[\mathcal{X}]$  denote the set of all polynomials.

Let  $f = \alpha_1 X_1 \alpha_2 \dots X_k \alpha_{k+1}$  be a monomial and let  $v$  be a vector. The *evaluation of  $f$  at  $v$* , denoted by  $f(v)$ , is the product  $\alpha_1 v_{X_1} \alpha_2 \dots \alpha_k v_{X_k} \alpha_{k+1}$ . We extend this to any polynomial: if  $f = \sum_{i=1}^k m_i$ , then  $f(v) = \sum_{i=1}^k m_i(v)$ .

A *system of polynomials* or polynomial system is a map  $\mathbf{f} : \mathcal{X} \rightarrow \mathcal{S}[\mathcal{X}]$ . We write  $\mathbf{f}_X$  for  $\mathbf{f}(X)$ . Every polynomial system induces a map from  $V$  to  $V$  by componentwise evaluation of the polynomials:  $\mathbf{f}(v)_X := \mathbf{f}_X(v)$  for all  $v \in V$ , and  $X \in \mathcal{X}$ . The following proposition, which follows easily from Kleene's theorem and the fact that  $\mathbf{f}$  is a monotone and continuous mapping, shows that any polynomial system  $\mathbf{f}$  has a least fixed-point  $\mu\mathbf{f}$ , which is by definition the least solution of  $\mathbf{X} = \mathbf{f}(\mathbf{X})$ .


**Proposition 1.** *A polynomial system  $\mathbf{f}$  has a unique least fixed-point  $\mu\mathbf{f}$ , i.e.,  $\mu\mathbf{f} = \mathbf{f}(\mu\mathbf{f})$ , and  $\mu\mathbf{f} \sqsubseteq v$  holds for all  $v$  with  $v = \mathbf{f}(v)$ . Further,  $\mu\mathbf{f}$  is the supremum (w.r.t.  $\sqsubseteq$ ) of the Kleene sequence  $(\mathbf{f}^i(\mathbf{0}))_{i \in \mathbb{N}}$ , where  $\mathbf{f}^i$  denotes the  $i$ -fold application of  $\mathbf{f}$ .*

### 3 Derivation Trees

We generalize the notion of derivation tree, as known from formal languages and grammars. We identify a node  $u$  of a (ordered) tree  $t$  with the subtree of  $t$  rooted at  $u$ . In particular, we identify a tree with its root.

Let  $\mathbf{f}$  be a polynomial system over a set  $\mathcal{X}$  of variables. A *derivation tree  $t$  of  $\mathbf{f}$*  is an ordered (finite) tree whose nodes are labelled with both a variable  $X$  and a monomial  $m$  of  $\mathbf{f}_X$ . We write  $\lambda_v$ , resp.  $\lambda_m$  for the corresponding labelling-functions. Moreover, if the monomial labelling of a node  $u$  is  $\lambda_m(u) = a_1 X_1 a_2 \dots X_s a_{s+1}$  for some  $s \geq 0$ , then  $u$  has exactly  $s$  children  $u_1, \dots, u_s$ , ordered from left to right, with  $\lambda_v(u_i) = X_i$  for all  $i = 1, \dots, s$ . A derivation tree  $t$  is an  *$X$ -tree* if  $\lambda_v(t) = X$ . The set of all  $X$ -trees of  $\mathbf{f}$  is denoted by  $\mathcal{T}_{\mathbf{f}, X}$ , or just by  $\mathcal{T}_X$  if  $\mathbf{f}$  is clear from the context.

The left part of Figure 1 shows a derivation tree of the system  $\mathbf{f}$  over the variables  $X$  and  $Y$  given by  $\mathbf{f}_X = aXYb + c$  and  $\mathbf{f}_Y = dX + Ye$ . The derivation trees of  $\mathbf{f}$  are very similar to the derivation trees of the context-free grammar with productions  $X \rightarrow aXYb|c$  and  $Y \rightarrow dX|Ye$ . For technical reasons, the nodes of “our” trees are labeled by “productions” (for instance, the label  $(X, aXYb)$  corresponds to the production  $X \rightarrow aXYb$ ). On the right of Figure 1 we show how the tree would look like according to the standard definition. The height  $h(t)$  of a derivation tree  $t$  is the length of a longest



**Fig. 1.** A derivation tree on the left, and its standard representation on the right

path from the root to a leaf. The set of  $X$ -trees (of  $\mathbf{f}$ ) of height *at most*  $h$  is denoted by  $\mathcal{T}_X^{(h)}$ . The yield  $\Upsilon(t)$  of a derivation tree  $t$  with  $\lambda_m(t) = a_1X_1a_2 \cdots X_s a_{s+1}$  is inductively defined to be  $\Upsilon(t) = a_1\Upsilon(t_1)a_2 \cdots \Upsilon(t_s)a_{s+1}$ . We extend the definition of  $\Upsilon$  to sets  $T \subseteq \mathcal{T}_X$  by setting  $\Upsilon(T) := \sum_{t \in T} \Upsilon(t)$ . E.g., the system  $\mathbf{f}$  defined above has exactly two  $X$ -trees of height at most 2: the tree consisting of a single node labeled by  $(X, c)$ , and the left tree of Figure 1. Their yields are  $c$  and  $acdc b$ , respectively, and so  $\Upsilon(\mathcal{T}_X^{(2)}) = c + acdc b$ . It follows  $\Upsilon(\mathcal{T}_X^{(2)}) = \mathbf{f}^3(\mathbf{0})_X$ , i.e., the yield of the  $X$ -trees of height at most 2 is equal to the “Kleene approximant”  $\mathbf{f}^3(\mathbf{0})_X$  from Proposition 1. The following proposition, easy to prove [4], shows that this is not a coincidence.

**Proposition 2.** For all  $h \in \mathbb{N}$  and  $X \in \mathcal{X}$ , we have  $\Upsilon(\mathcal{T}_X^{(h)}) = (\mathbf{f}^{h+1}(\mathbf{0}))_X$ .

Together with Proposition 1 we get:

**Corollary 1.**  $\mu\mathbf{f}_X = \Upsilon(\mathcal{T}_X)$ .

### 3.1 Derivation Tree Analysis

We say that a set  $T_X$  of  $X$ -trees satisfies the *embedding property* if  $\Upsilon(t) \sqsubseteq \Upsilon(T_X)$  holds for every  $X$ -tree  $t$ . Loosely speaking, the yield of every  $X$ -tree can be “embedded” in the yield of  $T_X$ . As addition is idempotent, the embedding property immediately implies that  $\Upsilon(T_X) \sqsubseteq \Upsilon(T_X)$ . Of course, as  $T_X \subseteq \mathcal{T}_X$ , we also have the other direction, which leads to the following result.


**Proposition 3.** Let  $\mathbf{f}$  be a system of polynomials over an io-semiring, and let  $X$  be a variable of  $\mathbf{f}$ . If a set  $T_X$  of  $X$ -trees of  $\mathbf{f}$  satisfies the embedding property, then  $\mu\mathbf{f} = \Upsilon(T_X)$ .

This proposition suggests a technique for the design of efficient algorithms computing  $\mu f$ : (1) define a set  $T_X$  of derivation trees whose yield is “easy to compute” in some io-semiring, and (2) identify “relevant” classes of io-semirings for which  $T_X$  satisfies the embedding property. By Proposition 3,  $\mu f$  is “easy to compute” for these classes. We call this technique *derivation tree analysis*.

## 4 Bamboos and their Yield

The difficulty of derivation tree analysis lies in finding a set  $T_X$  exhibiting a good balance between the contradictory requirements “easy to compute” and “relevant”: if  $T_X = \emptyset$  then the yield is trivial to compute, but  $T_X$  does not satisfy the embedding property in any interesting case. Conversely,  $T_X = \mathcal{T}_X$  trivially satisfies the embedding property for every io-semiring, but is not easy to compute. The main contribution of this paper is the identification of a class of derivation trees, *bamboos*, exhibiting this balance. In this section we define bamboos and show that their yield is the least solution of a system of *linear* equations easily derivable from  $f$ . The “easy to compute” part is justified by the fact that in most semirings used in practice linear equations are far easier to solve than polynomial equations (e.g. in the real semiring or the language semiring with union and concatenation as operations). The “relevance” of bamboos is justified in the next three sections.

**Definition 1.** Let  $f$  be a system of polynomials. A tree  $t \in \mathcal{T}_{f,X}$  is an  $X$ -bamboo if there is a path leading from the root of  $t$  to some leaf of  $t$ , the stem, such that the height of every subtree of  $t$  not containing a node of the stem is at most  $n - 1$ . The set of all  $X$ -bamboos of  $f$  is denoted by  $\mathcal{B}_{f,X}$ , or just by  $\mathcal{B}_X$  if  $f$  is clear from the context.



**Fig. 2.** A bamboo with its stem printed bold; on the right it is shown with its stem straightened.

In order to define the system of linear equations mentioned above we need the notion of differential of a system of polynomials.

**Definition 2.** Let  $f \in \mathcal{S}[\mathcal{X}]$  be a polynomial and let  $v \in V$  be a vector. The differential of  $f$  at  $v$  w.r.t. a variable  $X$  is the map  $D_X f|_v : V \rightarrow S$  inductively defined as follows:

$$D_X f|_v(\mathbf{a}) = \begin{cases} 0 & \text{if } f \in S \text{ or } f \in \mathcal{X} \setminus \{X\} \\ \mathbf{a}_X & \text{if } f = X \\ D_X g|_v(\mathbf{a}) \cdot h(v) + g(v) \cdot D_X h|_v(\mathbf{a}) & \text{if } f = g \cdot h \\ \sum_{i=1}^k D_X m_i|_v(\mathbf{a}) & \text{if } f = \sum_{i=1}^k m_i. \end{cases}$$

Further, we define the differential of  $f$  at  $\mathbf{v}$  by  $Df|_{\mathbf{v}}(\mathbf{a}) := \sum_{X \in \mathcal{X}} D_X f|_{\mathbf{v}}(\mathbf{a})$ . The differential of a system of polynomials  $\mathbf{f}$  at  $\mathbf{v}$  is defined componentwise by  $(D\mathbf{f}|_{\mathbf{v}}(\mathbf{a}))_X := D(\mathbf{f}_X)|_{\mathbf{v}}(\mathbf{a})$  for all  $X \in \mathcal{X}$ .

*Example 1.* For  $f(X, Y) = a \cdot X \cdot X \cdot Y \cdot b$ ,  $\mathbf{v} = (v_X, v_Y)$ ,  $\mathbf{c} = (c_X, c_Y)$  we have:

$$\begin{aligned} D_X f|_{\mathbf{v}}(\mathbf{c}) &= a \cdot c_X \cdot v_X \cdot v_Y \cdot b + a \cdot v_X \cdot c_X \cdot v_Y \cdot b \\ D_Y f|_{\mathbf{v}}(\mathbf{c}) &= a \cdot v_X \cdot v_X \cdot c_Y \cdot b \end{aligned}$$

Using differentials we define a particular linearization of a polynomial system.

**Definition 3.** Let  $\mathbf{f}$  be a system of  $n$  polynomials. The bamboo system  $\mathbf{f}_{\mathcal{B}}$  associated to  $\mathbf{f}$  is the linear system  $\mathbf{f}_{\mathcal{B}}(\mathbf{X}) = D\mathbf{f}|_{\mathbf{f}(\mathbf{0})}(\mathbf{X}) + \mathbf{f}(\mathbf{0})$ . The least solution of the system of equations  $\mathbf{X} = \mathbf{f}_{\mathcal{B}}(\mathbf{X})$  is denoted by  $\mu\mathbf{f}_{\mathcal{B}}$ .

Now we can state the relation between bamboos and bamboo systems.

**Theorem 1.** Let  $\mathbf{f}$  be a system of polynomials over an io-semiring. For every variable  $X$  of  $\mathbf{f}$  we have  $Y(\mathcal{B}_X) = (\mu\mathbf{f}_{\mathcal{B}})_X$ , i.e., the yield of the  $X$ -bamboos is equal to the  $X$ -component of the least solution of the bamboo system.

Together with Proposition 3 we get the following corollary.

**Corollary 2 (derivation tree analysis for bamboos).** Let  $\mathbf{f}$  be a system of polynomials over an io-semiring. If  $\mathcal{B}_X$  satisfies the embedding property for all  $X$ , i.e., for all  $X$ -trees  $t$  it holds  $Y(t) \sqsubseteq Y(\mathcal{B}_X)$ , then  $\mu\mathbf{f} = \mu\mathbf{f}_{\mathcal{B}}$ .

## 5 Star-Distributive Semirings

**Definition 4.** A commutative (w.r.t. multiplication) io-semiring  $\mathcal{S}$  is star-distributive if  $(a + b)^* = a^* + b^*$  holds for all  $a, b \in \mathcal{S}$ .

A commutative io-semiring is star-distributive whenever the natural order  $\sqsubseteq$  is total:

**Proposition 4.** Any totally ordered commutative io-semiring is star-distributive.

*Proof.* Let w.l.o.g.  $a \sqsubseteq b$ . Then  $(a + b)^* = b^* \sqsubseteq a^* + b^* \sqsubseteq (a + b)^*$ .  $\square$

In particular, the  $(\min, +)$ -semiring over the integers or reals is star-distributive.

We have already considered commutative idempotent semirings in [5] where we showed that  $\mu\mathbf{f}$  can be computed by solving  $n$  linear equation systems by means of a Newton-like method, improving the  $\mathcal{O}(3^n)$  bound of Hopkins and Kozen [11]. In this section we improve this result even further for star-distributive semirings: One single linear system, the bamboo system  $\mathbf{f}_{\mathcal{B}}$ , needs to be solved. This leads to an efficient algorithm for computing  $\mu\mathbf{f}$  in arbitrary star-distributive semirings. In Section 5.1 we instantiate this algorithm for the  $(\min, +)$ -semiring; in Section 5.2 we use it to improve the algorithm of [2] for computing the throughput of a context-free grammar.

We start by stating two useful properties of star-distributive semirings.


**Proposition 5.** In any star-distributive semiring the following equations hold:

(1)  $a^*b^* = a^* + b^*$ , and (2)  $(ab^*)^* = a^* + ab^*$ .

We can now state and prove our result:

**Theorem 2.**  $\mu f = \mu f_{\mathcal{B}}$  holds for polynomial systems  $f$  over star-distributive semirings.

*Proof Sketch (see the appendix for a complete proof).* The proof is by derivation tree analysis. So it suffices to discharge the precondition of Corollary 2. More precisely we show for any  $X$ -tree  $t$  that  $Y(t) \sqsubseteq Y(\mathcal{B}_X)$  holds. It suffices to consider the case where  $t$  is not an  $X$ -bamboo. Then the height of  $t$  is at least  $n$ , and so  $t$  is “pumpable”, i.e., one can choose a path  $p$  in  $t$  from the root to a leaf such that two different nodes on the path share the same variable-label. So  $t$  can be decomposed into three (partial) trees with yields  $a, b, c$ , respectively, such that  $Y(t) = abc$ , see the left side of Figure 3(a). Notice that, by commutativity of product,  $ab^*c$  is the yield of a set of trees obtained by



**Fig. 3.** “Unpumping” trees to make them bamboos

“pumping”  $t$ . We show  $ab^*c \sqsubseteq Y(\mathcal{B}_X)$  which implies  $Y(t) \sqsubseteq Y(\mathcal{B}_X)$ . As  $t$  is not an  $X$ -bamboo,  $t$  has a pumpable subtree disjoint from  $p$ . In this sketch we assume that it is a subtree of that part of  $t$  whose yield is  $a$ , see the right side of Figure 3(a). Now we have  $a = a_1a_2a_3$ , and so  $ab^*c = a_1a_2a_3b^*c \sqsubseteq a_1a_2^*a_3b^*c = a_1a_3b^*c + a_1a_2^*a_3c$ , where we used commutativity and Proposition 5(1) in the last step. Both summands in above sum are yields of sets of trees obtained by pumping pumpable trees smaller than  $t$ , see Figure 3(b). By an inductive argument those yields are both included in  $Y(\mathcal{B}_X)$ .  $\square$

### 5.1 The $(\min, +)$ -Semiring

Consider the “tropical” semiring  $\mathcal{R} = (\mathbb{R} \cup \{-\infty, \infty\}, \wedge, \text{+}_{\mathbb{R}}, \infty, 0)$ . By  $\wedge$  resp.  $\text{+}_{\mathbb{R}}$  we mean minimum resp. addition over the reals. Observe that the natural order  $\sqsubseteq$  is the order  $\geq$  on the reals.<sup>1</sup> As  $\mathcal{R}$  is totally ordered, Proposition 4 implies that  $\mathcal{R}$  is star-distributive. Assume for the rest of this section that  $f$  is a polynomial system over  $\mathcal{R}$  of degree at most 2. We can apply Theorem 2, i.e.,  $\mu f = \mu f_{\mathcal{B}}$  holds. This immediately suggests a polynomial algorithm to compute the least fixed-point: Compute  $f^n(\infty)$  by

<sup>1</sup> By symmetry, we could equivalently consider maximum instead of minimum.

performing  $n$  Kleene iterations, and solve the linear system  $\mathbf{X} = D\mathbf{f}|_{f^n(\infty)}(\mathbf{X}) \wedge \mathbf{f}(\infty)$ . The latter can be done by means of the Bellman-Ford algorithm.


*Example 2.* Consider the following equation system.

$$(X, Y, Z) = (-2 \wedge (Y \uparrow_{\mathbb{R}} Z), Z \uparrow_{\mathbb{R}} 1, X \wedge Y) =: \mathbf{f}(\mathbf{X})$$

We have  $\mathbf{f}(\infty) = (-2, \infty, \infty)$ ,  $\mathbf{f}^2(\infty) = (-2, \infty, -2)$ ,  $\mathbf{f}^3(\infty) = (-2, -1, -2)$ . The linear system  $\mathbf{X} = D\mathbf{f}|_{f^n(\infty)}(\mathbf{X}) \wedge \mathbf{f}(\infty) = \mathbf{f}_{\mathcal{B}}(\mathbf{X})$  looks as follows:

$$(X, Y, Z) = (-2 \wedge (-1 \uparrow_{\mathbb{R}} Z) \wedge (Y \uparrow_{\mathbb{R}} -2), Z \uparrow_{\mathbb{R}} 1, X \wedge Y).$$

This equation system corresponds in a straightforward way to the following graph.



We claim that the  $V$ -component of  $\mu\mathbf{f}_{\mathcal{B}}$  equals the least weight of any path from  $S$  to  $V$  where  $V \in \{X, Y, Z\}$ . To see this, notice that  $(\mathbf{f}_{\mathcal{B}}^k(\infty))_V$  corresponds to the least weight of any path from  $S$  to  $V$  of length at most  $k$ . The claim follows by Kleene's theorem. So we can compute  $\mu\mathbf{f}_{\mathcal{B}}$  with the Bellman-Ford algorithm. In our example,  $X, Y, Z$  are all reachable from  $S$  via a negative cycle, so  $\mu\mathbf{f}_{\mathcal{B}} = (-\infty, -\infty, -\infty)$ . By Theorem 2,  $\mu\mathbf{f} = \mu\mathbf{f}_{\mathcal{B}} = (-\infty, -\infty, -\infty)$ .  $\square$

The Bellman-Ford algorithm can be used here as it handles negative cycles correctly. The overall runtime of our algorithm to compute  $\mu\mathbf{f}$  is dominated by the Bellman-Ford algorithm. Its runtime is in  $\mathcal{O}(n \cdot m)$ , where  $m$  is the number of monomials appearing in  $\mathbf{f}$ . We conclude that our algorithm has the same asymptotic complexity as the “generalized Bellman-Ford” algorithm of [8]. It is by a factor of  $n$  faster than the algorithm deducible from [5] because our new algorithm uses the Bellman-Ford algorithm only once instead of  $n$  times.

## 5.2 Throughput of Grammars

In [2], a polynomial algorithm for computing the *throughput* of a context-free grammar was given. Now we show that the algorithm can be both simplified and accelerated by computing least fixed-points according to Theorem 2.

Let us define the problem following [2]. Let  $\Sigma$  be a finite alphabet and  $\rho : \Sigma \rightarrow \mathbb{N}$  a weight function. We extend  $\rho$  to words  $a_1 \cdots a_k \in \Sigma^*$  by setting  $\rho(a_1 \cdots a_k) := \rho(a_1) + \dots + \rho(a_k)$ .<sup>2</sup> The mean weight of a non-empty word  $w$  is defined as  $\bar{\rho}(w) := \rho(w)/|w|$ . The throughput of a non-empty language  $L \subseteq \Sigma^+$  is defined as the infimum of the mean weights of the words in  $L$ :  $tp(L) := \inf\{\bar{\rho}(w) \mid w \in L\}$ . Let  $G = (\Sigma, \mathcal{X}, P, S)$  be a context-free grammar and  $L = L(G)$  its language. The problem is to

<sup>2</sup> We write  $+$  for the addition of reals in this section.



compute  $tp(L)$ . As in [2] we assume that  $G$  has at most 2 symbols on the right hand side of every production and that  $L$  is non-empty and contains only non-empty words.

Note that we cannot simply construct a polynomial system having  $tp(L)$  as its least fixed-point, as the throughput of two non-terminals is not additive. In [2] an ingenious algorithm is proposed to avoid this problem. Assume we already know a routine, the *comparing routine*, that decides for a given  $t \in \mathbb{Q}$  whether  $tp(L) \geq t$  holds. Assume further that this routine has  $\mathcal{O}(N^k)$  time complexity for some  $k$ . Using the comparing routine we can approximate  $tp(L)$  up to any given accuracy by means of binary search. Let  $d = \max_{a \in \Sigma} \rho(a) - \min_{a \in \Sigma} \rho(a)$ . A dichotomy result of [2] shows that  $\mathcal{O}(N + \log d)$  iterations of binary search suffice to approximate  $tp(L)$  up to an  $\varepsilon$  that allows to compute the exact value of  $tp(L)$  in time  $\mathcal{O}(N^3)$ . This is proved by showing that, once a value  $t$  has been determined such that  $t - \varepsilon < tp(L) \leq t$ , one can:

- transform  $G$  in  $\mathcal{O}(N^3)$  time into a grammar  $G'$  of size  $\mathcal{O}(N^3)$  generating a finite language, and having the same throughput as  $G$  (this construction does not yet depend on  $tp(L)$ );
- compute the throughput of  $G'$  in linear time in the size of  $G'$ , i.e., in  $\mathcal{O}(N^3)$  time.

The full algorithm for the throughput runs then in  $\mathcal{O}(N^k(N + \log d)) + \mathcal{O}(N^3)$  time.

The algorithm of [2] and our new algorithm differ in the comparing routine. In the routine of [2] the transformation of  $G$  into the grammar  $G'$  is done *before*  $tp(L)$  has been determined. Then a linear time algorithm can be applied to  $G'$  to decide whether  $tp(L) \geq t$  holds. (This algorithm does not work for arbitrary context-free grammars, and that is why one needs to transform  $G$  into  $G'$ .) Since  $G'$  has size  $\mathcal{O}(N^3)$ , the comparing routine has  $k = 3$ , and so the full algorithm runs in  $\mathcal{O}(N^4 + N^3 \log d)$  time.

We give a more efficient comparing routine with  $k = 2$ . Given a  $t \in \mathbb{Q}$ , assign to each word  $w \in \Sigma^+$  its *throughput balance*  $\sigma_t(w) = \rho(w) - |w| \cdot t$ . Notice that  $\sigma_t(w) \geq 0$  if and only if  $\bar{\rho}(w) \geq t$ . Further, for two words  $w, u$  we now have  $\sigma_t(wu) = \sigma_t(w) + \sigma_t(u)$ . So we can set up a polynomial system  $\mathbf{X} = \mathbf{f}(\mathbf{X})$  over the tropical semiring  $\mathcal{R}$  where  $\mathbf{f}$  is constructed such that each variable  $X \in \mathcal{X}$  in the equation system corresponds to the minimum (infimum) throughput balance of the words derivable from  $X$ . More formally, define a map  $m$  by setting  $m(a) = \rho(a) - t$  for  $a \in \Sigma$  and  $m(X) = X$  for  $X \in \mathcal{X}$ . Extend  $m$  to words in  $(\Sigma \cup \mathcal{X})^*$  by setting  $m(\alpha_1 \cdots \alpha_k) = m(\alpha_1) + \cdots + m(\alpha_k)$ . Let  $P_X$  be the productions of  $G$  with  $X$  on the left hand side. Then set  $\mathbf{f}_X(\mathbf{X}) := \bigwedge_{(X \rightarrow w) \in P_X} m(w)$ . For instance, if  $P_X$  consists of the rules  $X \rightarrow aXY$  and  $X \rightarrow bZ$ , we have  $\mathbf{f}_X(\mathbf{X}) = \rho(a) - t + X + Y \wedge \rho(b) - t + Z$ .

It is easy to see that the relevant solution of the system  $\mathbf{X} = \mathbf{f}(\mathbf{X})$  is the least one w.r.t.  $\sqsubseteq$ , i.e.,  $(\mu \mathbf{f})_S \geq 0$  if and only if  $tp(L) \geq t$ . So we can use the algorithm from Section 5.1 as our comparing routine. This takes time  $\mathcal{O}(N^2)$  where  $N$  is the size of the grammar. With that comparing routine we obtain an algorithm for computing the throughput with  $\mathcal{O}(N^3 + N^2 \log d)$  runtime.

## 6 Lossy Semirings

**Definition 5.** An io-semiring  $\mathcal{S}$  is called *lossy* if  $1 \sqsubseteq a$  holds for all  $a \neq 0$ .

Note that by definition of natural order the requirement  $1 \sqsubseteq a$  is equivalent to  $a = a + 1$ . In the free semiring generated by a finite alphabet  $\Sigma$ , and augmented by the equation  $a = a + 1$  ( $a \in S \setminus \{0\}$ ), every language  $L \subseteq \Sigma^*$  is “downward closed”, i.e. for every word  $w = a_1 a_2 \dots a_l \in L$  all possible subwords  $\{a'_1 a'_2 \dots a'_l \mid a'_i \in \{\varepsilon, a_i\}\}$  are also included in  $L$ . By virtue of Higman’s lemma [10] the downward-closure of a context-free language is regular. This has been used in [1] for an efficient analysis of systems with unbounded, lossy FIFO channels. Downward closure was used there to model the loss of messages due to transmission errors.

We say that a system  $\mathbf{f}$  of polynomials is *clean* if  $\mu \mathbf{f}_X \neq 0$  for all  $X \in \mathcal{X}$ . Every system can be *cleaned* in linear time by removing the equations of all variables  $X$  such that  $\mu \mathbf{f}_X = 0$  and setting these variables to 0 in the other equations (the procedure is similar to the one that eliminates non-productive variables in context-free grammars). We consider only clean systems, and introduce a normal form for them.

**Definition 6.** Let  $\mathbf{f} \in \mathcal{S}[\mathcal{X}]^{\mathcal{X}}$  be a system of polynomials over a lossy semiring.  $\mathbf{f}$  is in quadratic normal form if every polynomial  $\mathbf{f}_X$  has the form

$$c + \sum_{Y, Z \in \mathcal{X}} a_{Y, Z} \cdot Y \cdot Z + \sum_{Y \in \mathcal{X}} b_{l, Y} \cdot Y \cdot b_{r, Y}$$

where (i)  $c \in S \setminus \{0\}$ , (ii)  $a_{Y, Z} \in \{0, 1\}$ , and (iii) if  $\sum_{Z \in \mathcal{X}} a_{Y, Z} \neq 0$ , then  $b_{l, Y} \neq 0 \neq b_{r, Y}$  for all  $Y, Z \in \mathcal{X}$ .

**Lemma 1.** For every clean  $\mathbf{g} \in \mathcal{S}[\mathcal{X}]^{\mathcal{X}}$  we can construct in linear time a system  $\mathbf{f} \in \mathcal{S}[\mathcal{X}']^{\mathcal{X}'}$  in quadratic normal form, where  $\mathcal{X} \subseteq \mathcal{X}'$  and  $\mu \mathbf{g}_X = \mu \mathbf{f}_X$  for all  $X \in \mathcal{X}$ .


*Proof Sketch.* Note that, as  $\mathbf{g}$  is clean, we have  $1 \sqsubseteq \mu \mathbf{g}$ . Hence, requirement (i) is no restriction. The transformation that normalizes a system is similar to the one that brings a context-free grammar into Chomsky normal-form (CNF). The superset  $\mathcal{X}' \supset \mathcal{X}$  results from the introduction of new variables by this transformation into CNF.  $\square$

Our main result in this section is that for *strongly connected* systems  $\mathbf{f}$  in quadratic normal form we again have that  $\mu \mathbf{f} = \mu \mathbf{f}_{\mathcal{B}}$ . We then show how this result leads to an algorithm for arbitrary systems.

Given two variables  $X, Y \in \mathcal{X}$ , we say that  $X$  depends on  $Y$  (w.r.t.  $\mathbf{f}$ ) if  $Y$  occurs in a monomial of  $\mathbf{f}_X$  or there is a variable  $Z$  such that  $X$  depends on  $Z$  and  $Z$  depends on  $Y$ . The system  $\mathbf{f}$  is *strongly connected* if  $X$  depends on  $Y$  for all variables  $X, Y$ .

**Theorem 3.**  $\mu \mathbf{f} = \mu \mathbf{f}_{\mathcal{B}}$  holds for strongly connected polynomial systems  $\mathbf{f}$  in quadratic normal form over lossy semirings.

*Proof Sketch.* We consider a concrete example of a tree  $t$  that is not a bamboo, and show how to construct a bamboo  $\hat{t}$  such that  $Y(t) \sqsubseteq Y(\hat{t})$ . The general procedure for all non-bamboos can be found in the appendix. Let  $\mathcal{X} = \{X, Y\}$ , and  $\mathbf{f}$  with  $\mathbf{f}_X = XY + X + Y + a$ , and  $\mathbf{f}_Y = X + Y + b$ . Consider the  $X$ -tree  $t$  depicted on the left of the picture below, where  $t_r$  is some bamboo of height at least 2 (we inductively assume that the original subtree has already been replaced by a bamboo with at least the same yield). Since the left subtree of  $t$  has height 2,  $t$  itself is not a bamboo. Let  $s$  denote the



left-most leaf of  $t$ , and let  $r$  be the parent of  $s$ . In our example, we assume that  $r$  has  $s$  as its only child. Then we proceed as follows:

(i) We remove from  $t$  the leaf  $s$ , and turn its father  $r$  into a leaf. Here, we make use of the assumption that  $\mathbf{f}$  is in quadratic normal form, and so every polynomial of  $\mathbf{f}$  contains a constant monomial, in our example  $b$ . We change the monomial-label of  $r$  to  $b$ , and obtain the tree  $t'$ , which is a derivation tree of  $\mathbf{f}$ . Moreover,  $t'$  is a bamboo, because its left subtree has now height 1, and its right subtree  $t_r$  is a bamboo.

(ii) We prepend a (partial) derivation tree on top of  $t_r$  having two linear chains as subtrees: the left chain leads to the leaf  $s$ , and the right chain leads to  $t'$ . This gives us the tree  $\hat{t}$  depicted on the right of the picture above. The proof of Theorem 3 shows that these chains exist and have at most length  $n - 1$  (in our example  $n - 1 = 1$ ). It follows that  $\hat{t}$  is a bamboo itself with stem  $\hat{t} - t'' - t' - t_r$ , and so  $\Upsilon(\hat{t}) \sqsubseteq \Upsilon(\mathcal{B}_X)$ .

We have  $\Upsilon(t) = a \cdot \Upsilon(t_r)$  and  $\Upsilon(\hat{t}) = a \cdot b \cdot \Upsilon(t_r)$ . Since the semiring is lossy, we have  $1 \sqsubseteq b$  and so  $\Upsilon(t) \sqsubseteq \Upsilon(\hat{t})$ . Notice that, since product is not necessarily commutative, it is important that  $a$  is the first factor of both yields.  $\square$

Because of the preceding theorem, given a strongly connected system  $\mathbf{f}$ , we may use the linear system  $\mathbf{f}_{\mathcal{B}}(\mathbf{X}) = \mathbf{f}(\mathbf{0}) + D\mathbf{f}|_{\mathbf{f}^n(\mathbf{0})}(\mathbf{X})$  for calculating  $\mu\mathbf{f}$ . As  $\mathbf{f}$  is strongly connected,  $\mathbf{f}_{\mathcal{B}}$  is also strongly connected. The least fixed-point of such a strongly connected linear system  $\mathbf{f}_{\mathcal{B}}$  is easily calculated: all non-constant monomials appearing in  $\mathbf{f}_{\mathcal{B}}$  have the form  $b_l X b_r$  for some  $X \in \mathcal{X}$ , and  $b_l, b_r \in S \setminus \{0\}$ . As  $\mathbf{f}_{\mathcal{B}}$  is strongly connected, every polynomial  $(\mathbf{f}_{\mathcal{B}})_Y$  is substituted for  $Y$  in  $(\mathbf{f}_{\mathcal{B}})_X$  again and again when calculating the Kleene sequence  $(\mathbf{f}_{\mathcal{B}}^k(\mathbf{0}))_{k \in \mathbb{N}}$ . So, let  $l$  be the sum of all left-handed coefficients  $b_l$  (appearing in any  $\mathbf{f}_X$ ), and similarly define  $r$ . We then have  $(\mu\mathbf{f}_{\mathcal{B}})_X = l^* (\sum_{Y \in \mathcal{X}} \mathbf{f}_Y(\mathbf{0})) r^*$  for all  $X \in \mathcal{X}$ .

If  $\mathbf{f}$  is not strongly connected, we first decompose  $\mathbf{f}$  into strongly connected subsystems, and then we solve these systems bottom-up. Note that substituting the solutions from underlying SCCs into a given SCC leads to a new system in normal form. As there are at most  $n = |\mathcal{X}|$  many strongly connected components for a given system  $\mathbf{f} \in \mathcal{S}[\mathcal{X}]^{\mathcal{X}}$ , we obtain the following theorem which was first stated explicitly for context-free grammars in [3].

**Theorem 4.** *The least fixed-point  $\mu\mathbf{f}$  of a polynomial system  $\mathbf{f}$  over a lossy semiring is representable by regular expressions over  $\mathcal{S}$ . If  $\mathbf{f}$  is in normal form  $\mu\mathbf{f}$  can be calculated solving at most  $n$  bamboo systems.*

## 7 1-bounded Semirings

**Definition 7.** An io-semiring  $\mathcal{S}$  is called 1-bounded if  $a \sqsubseteq 1$  holds for all  $a \in \mathcal{S}$ .

Natural examples are the tropical semiring over the natural numbers  $(\mathbb{N} \cup \{\infty\}, \wedge, +, \infty, 0)$  and the “maximum-probability” semiring  $([0, 1], \vee, \cdot, 0, 1)$ , where  $\wedge$  and  $\vee$  denote minimum and maximum, respectively. Notice that any commutative 1-bounded semiring is star-distributive (as  $a^* = 1$  for all  $a$ ), but not all 1-bounded semirings have commutative multiplication. Consider for example the semiring of those languages  $L$  over  $\Sigma$  that are *upward-closed*, i.e.,  $w \in L$  implies  $u \in L$  for all  $u$  such that  $w$  is a subword of  $u$ . This semiring is 1-bounded and has  $\Sigma^*$  as 1-element. Upward-closed languages form a natural dual to downward-closed languages from the previous section.

We show that  $\mu f$  can be computed very easily in the case of 1-bounded semirings:

**Theorem 5.**  $\mu f = f^n(\mathbf{0})$  holds for polynomial systems over 1-bounded semirings.

*Proof Sketch.* Recall that, by Proposition 2, we have  $(f^n(\mathbf{0}))_X = Y(\mathcal{T}_X^{(n-1)})$ , where  $\mathcal{T}_X^{(n-1)}$  contains all  $X$ -trees of height at most  $n - 1$ . We proceed by derivation tree analysis, i.e., by discharging the precondition of Proposition 3. So it suffices to show that for any  $X$ -tree  $t$  there is an  $X$ -tree  $t'$  of height at most  $n - 1$  with  $Y(t) \sqsubseteq Y(t')$ . Such a tree  $t'$  can be constructed by pruning  $t$  as long as some variable label occurs more than once along any path.  $\square$

Theorem 5 appears to be rather easy from our point of view, i.e., from the point of view of derivation trees. However, even this simple result has very concrete applications in the domain of interprocedural program analysis [13]. The main algorithms of [13], the so-called *post\** and *pre\** algorithms, can be seen as solvers of fixed-point equations over *bounded* semirings, which are semirings that do not have infinite ascending chains. Those solvers are based on Kleene’s iteration and the complexity result given there depends on the maximal length of ascending chains in the semiring (cf. [13], page 28). Such a bound may not exist, and does not exist for the tropical semiring over the natural numbers  $(\mathbb{N} \cup \{\infty\}, \wedge, +, \infty, 0)$  which is considered as an example in [13], pages 13 and 18. However, Theorem 5 can be applied to this semiring, which shows that the program analysis algorithms of [13] applied to 1-bounded semirings are polynomial-time algorithms, independent of the length of chains in the semiring.

## 8 Conclusion

We have shown that derivation tree analysis, a proof technique first introduced in [5], is an efficient tool for the design of efficient fixed-point algorithms on io-semirings. We have considered three classes of io-semirings with applications to language theory and verification. We have shown that for star-distributive semirings and lossy semirings the least fixed-point of a polynomial system of equations is equal to the least fixed-point of a *linear system*, the bamboo system. This improves the results of [5]: The generic algorithm given there requires to solve  $N$  different systems of linear equations in the star-distributive case (where  $N$  is the original number of polynomial equations), and is not applicable to the lossy case.

We have used our results to design an efficient fixed-point algorithm for the  $(\min, +)$ -semiring. In turn, we have applied this algorithm to provide a cubic algorithm for computing the throughput of a context-free language, improving the  $\mathcal{O}(N^4)$  upper bound obtained by Caucal et al. in [2].

For lossy semirings, derivation tree analysis based on bamboos has led to an algebraic generalization of a result of Courcelle stating that the downward-closure of a context-free language is effectively regular. Finally we have used derivation tree analysis to derive a simple proof that  $\mu f = f^n(\mathbf{0})$  holds for 1-bounded semirings, with some applications in interprocedural program analysis.

## References

1. P. A. Abdulla, A. Bouajjani, and B. Jonsson. On-the-fly analysis of systems with unbounded, lossy FIFO channels. In *CAV'98*, LNCS 1427, pages 305–318. Springer, 1998.
2. D. Caucal, J. Czyzowicz, W. Fraczak, and W. Rytter. Efficient computation of throughput values of context-free languages. In *CIAA'07*, LNCS 4783, pages 203–213. Springer, 2007.
3. B. Courcelle. On constructing obstruction sets of words. *EATCS Bulletin*, 44:178–185, 1991.
4. J. Esparza, S. Kiefer, and M. Luttenberger. An extension of Newton's method to  $\omega$ -continuous semirings. In *DLT'07*, LNCS 4588, pages 157–168. Springer, 2007.
5. J. Esparza, S. Kiefer, and M. Luttenberger. On fixed point equations over commutative semirings. In *STACS'07*, LNCS 4397, pages 296–307. Springer, 2007.
6. J. Esparza, A. Kučera, and R. Mayr. Model checking probabilistic pushdown automata. *Logical Methods in Computer Science*, 2006.
7. K. Etessami and M. Yannakakis. Recursive Markov chains, stochastic grammars, and monotone systems of nonlinear equations. In *STACS*, pages 340–352, 2005.
8. T. Gawlitza and H. Seidl. Precise fixpoint computation through strategy iteration. In *ESOP'07*, LNCS 4421, pages 300–315. Springer, 2007.
9. T.E. Harris. *The Theory of Branching Processes*. Springer, 1963.
10. G. Higman. Ordering by divisibility in abstract algebras. *Proc. London Math. Soc.*, 2, 1952.
11. M. W. Hopkins and D. Kozen. Parikh's theorem in commutative Kleene algebra. In *LICS'99*.
12. F. Nielson, H.R. Nielson, and C. Hankin. *Principles of Program Analysis*. Springer, 1999.
13. T. Reps, S. Schwoon, S. Jha, and D. Melski. Weighted pushdown systems and their application to interprocedural dataflow analysis. *Science of Computer Programming*, 58(1–2):206–263, October 2005. Special Issue on the Static Analysis Symposium 2003.

## Appendix

### A Proofs of Section 4

**Theorem 1.** *Let  $f$  be a system of polynomials over an io-semiring. For every variable  $X$  of  $f$  we have  $Y(\mathcal{B}_X) = (\mu f_{\mathcal{B}})_X$ , i.e., the yield of the  $X$ -bamboos is equal to the  $X$ -component of the least solution of the bamboo system.*

*Proof.* By definition, we have

$$f_{\mathcal{B}}(X) = f(0) + Df|_{f^n(0)}(X).$$

Its Kleene sequence, thus, becomes

$$\begin{aligned} f_{\mathcal{B}}^0(0) &= f(0) \\ f_{\mathcal{B}}^1(0) &= f(0) + Df|_{f^n(0)}(f(0)) \\ f_{\mathcal{B}}^2(0) &= f(0) + Df|_{f^n(0)}(f(0)) + \underbrace{Df|_{f^n(0)}(Df|_{f^n(0)}(f(0)))}_{=: Df|_{f^n(0)}^2(f(0))} \\ &\vdots \\ f_{\mathcal{B}}^k(0) &= f(0) + Df|_{f^n(0)}(f(0)) + \dots + Df|_{f^n(0)}^{k-1}(f(0)). \end{aligned}$$

As  $\mu f_{\mathcal{B}} = \sup f_{\mathcal{B}}^k(0)$ , we get

$$\mu f_{\mathcal{B}} = \sum_{k \in \mathbb{N}} Df|_{f^n(0)}^k(f(0)),$$

where  $Df|_{f^n(0)}^k$  denotes the  $k$ -fold application of  $Df|_{f^n(0)}$ . In particular, we have  $Df|_{f^n(0)}^0(f(0)) = f(0)$ .

Let  $\mathcal{B}_X^{(h)}$  be the set of  $X$ -bamboos (w.r.t.  $f$ ) of height *at most*  $h$ . Similarly,  $\mathcal{B}_X$  denotes the set of all bamboos of height *at most*  $h$ .

$\sqsupseteq$ -direction: We first prove that

$$Y(\mathcal{B}_X) \sqsupseteq (\mu f_{\mathcal{B}})_X.$$

It suffices to show (for  $h \geq 0$ ) that

$$Y(\mathcal{B}_X^{(h+n-1)}) \sqsupseteq \left( Df|_{f^n(0)}^h(f^n(0)) \right)_X$$

holds for all  $X \in \mathcal{X}$  as this immediately implies

$$\begin{aligned} Y(\mathcal{B}_X) &= \sum_{h \in \mathbb{N}} Y(\mathcal{B}_X^{(h+n-1)}) \\ &\sqsupseteq \sum_{h \in \mathbb{N}} \left( Df|_{f^n(0)}^h(f^n(0)) \right)_X \\ &\sqsupseteq \sum_{h \in \mathbb{N}} \left( Df|_{f^n(0)}^h(f(0)) \right)_X \\ &= (\mu f_{\mathcal{B}})_X. \end{aligned}$$

We proceed by induction on  $h$ : For  $h = 0$ , we have  $\mathcal{B}_X^{(n-1)} = \mathcal{T}_X^{(n-1)}$ , and  $\Upsilon(\mathcal{T}_X^{(n-1)}) = \mathbf{f}^n(\mathbf{0})_X$  (cf. prop. 2). Thus,

$$\Upsilon(\mathcal{B}_X^{(n-1)}) = \Upsilon(\mathcal{T}_X^{(n-1)}) = \mathbf{f}^n(\mathbf{0})_X = \left( D\mathbf{f}|_{\mathbf{f}^n(\mathbf{0})}(\mathbf{f}^n(\mathbf{0})) \right)_X$$

follows immediately.

Consider therefore  $\left( D\mathbf{f}|_{\mathbf{f}^n(\mathbf{0})}^{h+1}(\mathbf{f}^n(\mathbf{0})) \right)_X$  for  $h \geq 0$ , and let  $\Upsilon(\mathcal{B}^{(h)})$  denote the vector defined by  $\Upsilon(\mathcal{B}^{(h)})_X := \Upsilon(\mathcal{B}_X^{(h)})$ . We then have by induction on  $h$  that

$$\left( D\mathbf{f}|_{\mathbf{f}^n(\mathbf{0})}^{h+1}(\mathbf{f}^n(\mathbf{0})) \right)_X \sqsubseteq \left( D\mathbf{f}|_{\mathbf{f}^n(\mathbf{0})}(\Upsilon(\mathcal{B}^{(h+n-1)})) \right)_X$$

Assume that  $\mathbf{f}_X = \sum_{i=1}^k m_i$  where  $m_1, \dots, m_k$  are monomials. As addition is idempotent, we may assume these monomials are pairwise different. By definition, we have

$$\begin{aligned} \left( D\mathbf{f}|_{\mathbf{f}^n(\mathbf{0})}(\Upsilon(\mathcal{B}^{(h+n-1)})) \right)_X &= D\mathbf{f}_X|_{\mathbf{f}^n(\mathbf{0})}(\Upsilon(\mathcal{B}^{(h+n-1)})) \\ &= \sum_{i=1}^k Dm_i|_{\mathbf{f}^n(\mathbf{0})}(\Upsilon(\mathcal{B}^{(h+n-1)})). \end{aligned}$$

If all monomials  $m_1, \dots, m_k$  have degree 0, then  $D\mathbf{f}_X|_{\mathbf{v}}(\mathbf{a}) = 0$  for all  $\mathbf{v}, \mathbf{a} \in V$ . But this also implies that  $\mathcal{B}^{(h)} = \mathcal{T}_X^{(0)}$  also holds for all  $h \geq 1$ , hence, we may assume that there is at least one monomial of degree at least one. Let  $m \in \{m_1, \dots, m_k\}$  be such a monomial with  $m = a_1 X_1 a_2 \dots X_l a_{l+1}$  (for some  $l \geq 1$ ). We then have

$$Dm|_{\mathbf{f}^n(\mathbf{0})}(\Upsilon(\mathcal{B}^{(h+n-1)})) = \sum_{Y \in \mathcal{X}} D_Y m|_{\mathbf{f}^n(\mathbf{0})}(\Upsilon(\mathcal{B}^{(h+n-1)}))$$

by definition. Consider a  $Y \in \{X_1, \dots, X_l\}$ , i.e. a variable appearing in  $m$  (for the remaining variables, the differential is again 0), and let  $\text{pos}_Y(m) = \{i \mid X_i = Y\}$  be the set of ‘‘positions of  $Y$  in  $m$ ’’. We then may write

$$\begin{aligned} & D_Y m|_{\mathbf{f}^n(\mathbf{0})}(\Upsilon(\mathcal{B}^{(h+n-1)})) \\ &= \sum_{p \in \text{pos}_Y(m)} \left( \prod_{q=1}^{p-1} a_q \cdot \mathbf{f}^n(\mathbf{0})_{X_q} \right) \cdot a_p \cdot \Upsilon(\mathcal{B}^{(h+n-1)})_{X_p} \cdot \left( \prod_{q=p+1}^l a_q \cdot \mathbf{f}^n(\mathbf{0})_{X_q} \right) \cdot a_{l+1} \\ &= \sum_{p \in \text{pos}_Y(m)} \left( \prod_{q=1}^{p-1} a_q \cdot \mathbf{f}^n(\mathbf{0})_{X_q} \right) \cdot a_p \cdot \Upsilon(\mathcal{B}_{X_p}^{(h+n-1)}) \cdot \left( \prod_{q=p+1}^l a_q \cdot \mathbf{f}^n(\mathbf{0})_{X_q} \right) \cdot a_{l+1} \\ &= \sum_{p \in \text{pos}_Y(m)} \left( \prod_{q=1}^{p-1} a_q \cdot \Upsilon(\mathcal{T}^{(n-1)})_{X_q} \right) \cdot a_p \cdot \Upsilon(\mathcal{B}_{X_p}^{(h+n-1)}) \cdot \left( \prod_{q=p+1}^l a_q \cdot \Upsilon(\mathcal{T}^{(n-1)})_{X_q} \right) \cdot a_{l+1} \\ &= \sum_{p \in \text{pos}_Y(m)} \left( \prod_{q=1}^{p-1} a_q \cdot \Upsilon(\mathcal{T}_{X_q}^{(n-1)}) \right) \cdot a_p \cdot \Upsilon(\mathcal{B}_{X_p}^{(h+n-1)}) \cdot \left( \prod_{q=p+1}^l a_q \cdot \Upsilon(\mathcal{T}_{X_q}^{(n-1)}) \right) \cdot a_{l+1} \\ &= \sum_{p \in \text{pos}_Y(m)} \left( \prod_{q=1}^{p-1} a_q \cdot \sum_{t \in \mathcal{T}_{X_q}^{(n-1)}} \Upsilon(t) \right) \cdot a_p \cdot \left( \sum_{t \in \mathcal{B}_{X_p}^{(h+n-1)}} \Upsilon(t) \right) \cdot \left( \prod_{q=p+1}^l a_q \cdot \sum_{t \in \mathcal{T}_{X_q}^{(n-1)}} \Upsilon(t) \right) \cdot a_{l+1} \end{aligned}$$

But this last sum is simply the yield of all  $X$ -bamboos  $t \in \mathcal{B}_X^{(h+n)}$  with  $\lambda_2(t) = m$  having height at least 1, and at most  $h + n$ . As for  $t \in \mathcal{B}_X^{(h)}$  we have  $\lambda_2(t) \in \{m_1, \dots, m_k\}$ , we get by idempotent addition

$$\left( D\mathbf{f}|_{\mathbf{f}^n(\mathbf{0})}(\mathbf{Y}(\mathcal{B}^{(h+n-1)})) \right)_X \sqsubseteq \mathbf{Y}(\mathcal{B}_X^{(h+n)}).$$

$\sqsubseteq$ -direction: We now turn to the proof of

$$\mathbf{Y}(\mathcal{B}_X) \sqsubseteq (\mu\mathbf{f}_\mathcal{B})_X.$$

As addition is idempotent, it suffices to show that

$$\mathbf{Y}(t) \sqsubseteq (\mu\mathbf{f}_\mathcal{B})_X.$$

We proceed by induction on the number of nodes in  $t$ . If  $t$  has just one node then  $\mathbf{Y}(t) \sqsubseteq (\mathbf{f}(\mathbf{0}))_X \sqsubseteq (\mu\mathbf{f}_\mathcal{B})_X$ . For the induction step,  $t$  has children. So assume w.l.o.g. that  $\lambda_2(t) = a_1 X_1 \cdots X_s a_{s+1}$  for some  $s \geq 1$ . Denote the children of  $t$  by  $t_1, \dots, t_s$ . Furthermore we assume w.l.o.g. that the backbone of  $t$  goes through  $t_1$ . Hence,  $t_1$  is itself a bamboo having less nodes than  $t$ . By induction we have  $\mathbf{Y}(t_1) \sqsubseteq (\mu\mathbf{f}_\mathcal{B})_{X_1}$ . As  $t$  is a bamboo, the other children  $t_2, \dots, t_s$  have a height of at most  $n - 1$ . It is easy to see (cf. [4]) that this implies

$$\mathbf{Y}(t_r) \sqsubseteq (\mathbf{f}^n(\mathbf{0}))_{X_r} \text{ for all } 2 \leq r \leq s. \quad (1)$$

Now we have:

$$\begin{aligned} \mathbf{Y}(t) &= a_1 \mathbf{Y}(t_1) \cdots \mathbf{Y}(t_s) a_{s+1} && \text{(def. of yield } \mathbf{Y}) \\ &\sqsubseteq a_1 (\mu\mathbf{f}_\mathcal{B})_{X_1} a_2 \mathbf{Y}(t_2) \cdots \mathbf{Y}(t_s) a_{s+1} && \text{(by induction)} \\ &\sqsubseteq a_1 (\mu\mathbf{f}_\mathcal{B})_{X_1} a_2 (\mathbf{f}^n(\mathbf{0}))_{X_2} \cdots (\mathbf{f}^n(\mathbf{0}))_{X_s} a_{s+1} && \text{(Equation (1))} \\ &\sqsubseteq D_{X_1} (a_1 X_1 \cdots X_s a_{s+1})|_{\mathbf{f}^n(\mathbf{0})} (\mu\mathbf{f}_\mathcal{B}) && \text{(def. of differentials)} \\ &\sqsubseteq D_{X_1} \mathbf{f}_X|_{\mathbf{f}^n(\mathbf{0})} (\mu\mathbf{f}_\mathcal{B}) && (t \in \mathcal{B}_X) \\ &\sqsubseteq D\mathbf{f}_X|_{\mathbf{f}^n(\mathbf{0})} (\mu\mathbf{f}_\mathcal{B}) && \text{(def. of differentials)} \\ &= (D\mathbf{f}|_{\mathbf{f}^n(\mathbf{0})} (\mu\mathbf{f}_\mathcal{B}))_X && \text{(def. of differentials)} \\ &\sqsubseteq (\mu\mathbf{f}_\mathcal{B})_X \end{aligned}$$

□

## B Proofs of Section 5

**Proposition 5.** *In any star-distributive semiring, the following equations hold:*

- (1)  $a^* b^* = a^* + b^*$
- (2)  $(ab^*)^* = a^* + ab^*$

*Proof.*



- (1) The equation  $a^*b^* = (a + b)^*$  holds in any commutative idempotent semiring. By star-distributivity,  $(a + b)^* = a^* + b^*$ .
- (2) In any commutative io-semiring, we have  $(ab^*)^* = 1 + aa^*b^*$  (see e.g. [11]). By (1), we have  $1 + aa^*b^* = 1 + aa^* + ab^* = a^* + ab^*$ .  $\square$

**Theorem 2.**  $\mu f = \mu f_{\mathcal{B}}$  holds for polynomial systems  $f$  over star-distributive semirings.

*Proof.* We will need the following notation: If  $t'$  is a subtree of a derivation tree  $t$ , we write  $t = \widehat{t} \cdot t'$  where  $\widehat{t}$  is the partial derivation tree obtained from  $t$  by removing  $t'$ . If, in addition,  $t' = \widehat{t}' \cdot t''$ , and  $t'$  and  $t''$  have the same variable-label, we say the decomposition  $t = \widehat{t} \cdot \widehat{t}' \cdot t''$  is *pumpable*, because  $\widehat{t} \cdot (\widehat{t}')^i \cdot t''$  is a valid tree for all  $i \geq 0$ . We define  $\widehat{t} \cdot (\widehat{t}')^* \cdot t'' = \{\widehat{t} \cdot (\widehat{t}')^i \cdot t'' \mid i \geq 0\}$ . Notice that, due to commutativity of product, it holds  $Y(\widehat{t} \cdot (\widehat{t}')^* \cdot t'') = Y(\widehat{t}) \cdot Y(\widehat{t}')^* \cdot Y(t'')$ . We call this yield the *pumping yield* of the decomposition  $t = \widehat{t} \cdot \widehat{t}' \cdot t''$ .

The proof is by derivation tree analysis. So it suffices to discharge the precondition of Corollary 2. More precisely we need to show that, for any  $X$ -tree  $t$ , we have  $Y(t) \sqsubseteq Y(\mathcal{B}_X)$ . If  $t$  does not have a pumpable decomposition, then  $t$  has a height of at most  $n - 1$ , hence  $t \in \mathcal{B}_X$  and so  $Y(t) \sqsubseteq Y(\mathcal{B}_X)$ . It remains to show: if  $t$  has a pumpable decomposition  $t = \widehat{t} \cdot \widehat{t}_1 \cdot t'_1$ , then  $Y(t) \sqsubseteq \mathcal{B}_X$ . In fact, we show  $Y(\widehat{t} \cdot (\widehat{t}_1)^* \cdot t'_1) \sqsubseteq Y(\mathcal{B}_X)$ , which is stronger because  $Y(t) \sqsubseteq Y(\widehat{t} \cdot (\widehat{t}_1)^* \cdot t'_1)$ .

Denote by  $\#(t)$  the number of nodes in a tree  $t$ . We assign to a pumpable decomposition  $t = \widehat{t} \cdot \widehat{t}_1 \cdot t'_1$  a *size* by setting  $size(t = \widehat{t} \cdot \widehat{t}_1 \cdot t'_1) = (\#(t), \#(\widehat{t}_1 \cdot t'_1))$ . We define a total order on sizes by setting  $(i, j) \triangleleft (i', j')$  if either  $i < i'$  or  $i = i'$  and  $j < j'$ . We use this order to prove by induction that for any size  $(i, j)$ , if there is a pumpable decomposition  $t = \widehat{t} \cdot \widehat{t}_1 \cdot t'_1$  of size  $(i, j)$ , then  $Y(\widehat{t} \cdot (\widehat{t}_1)^* \cdot t'_1) \sqsubseteq Y(\mathcal{B}_X)$ .

The induction base is trivial because trees  $t$  with  $\#(t) = 1$  do not have a decomposition. For the induction step, let  $t$  be an  $X$ -tree and let  $t = \widehat{t} \cdot \widehat{t}_1 \cdot t'_1$  be pumpable. Choose a path  $p$  in  $t$  from the root to a leaf through  $t'_1$ . If  $p$  is a valid stem of an  $X$ -bamboo, then all trees in  $\widehat{t} \cdot (\widehat{t}_1)^* \cdot t'_1$  are  $X$ -bamboos, so  $Y(\widehat{t} \cdot (\widehat{t}_1)^* \cdot t'_1) \sqsubseteq \mathcal{B}_X$ . Hence, assume that  $p$  is not a valid stem, i.e., there is some subtree of  $t$ , disjoint from  $p$ , with height at least  $n$ . So this tree has a subtree  $t_2 = \widehat{t}_2 \cdot t'_2$  such that  $t_2$  and  $t'_2$  have the same variable-label. We distinguish two cases.

- (a) Let  $t_2$  *not* be a subtree of  $\widehat{t}_1$ . Then  $\widehat{t}_1$  and  $\widehat{t}_2$  are disjoint and so there exists a  $\widetilde{y}$  such that  $Y(t) = \widetilde{y} \cdot Y(\widehat{t}_1) \cdot Y(\widehat{t}_2)$ . Then:

$$\begin{aligned} Y(\widehat{t} \cdot (\widehat{t}_1)^* \cdot t'_1) &= \widetilde{y} \cdot Y(\widehat{t}_1)^* \cdot Y(\widehat{t}_2) \\ &\sqsubseteq \widetilde{y} \cdot Y(\widehat{t}_1)^* \cdot Y(\widehat{t}_2)^* \quad (\text{def. of Kleene } *) \\ &= \widetilde{y} \cdot Y(\widehat{t}_1)^* + \widetilde{y} \cdot Y(\widehat{t}_2)^* \quad (\text{Prop. 5 (1)}) \end{aligned}$$

The expression  $\widetilde{y} \cdot Y(\widehat{t}_1)^*$  equals the pumping yield of a decomposition of an  $X$ -tree which is obtained from  $t$  by removing the substructure  $\widehat{t}_2$ . Similarly, the expression  $\widetilde{y} \cdot Y(\widehat{t}_2)^*$  is equal to the pumping yield of a decomposition of an  $X$ -tree which is obtained from  $t$  by removing the substructure  $\widehat{t}_1$ . By induction on the size, both of those pumping yields are  $\sqsubseteq Y(\mathcal{B}_X)$ .

(b) Let  $t_2$  be a subtree of  $\widehat{t}_1$ . Then we can write  $\widehat{t}_1 = \widehat{\widehat{t}_1} \cdot \widehat{t}_2 \cdot t'_2$ . We have:

$$\begin{aligned}
Y(\widehat{t} \cdot (\widehat{t}_1)^* \cdot t'_1) &= Y(\widehat{t}) \cdot Y(\widehat{\widehat{t}_1} \cdot \widehat{t}_2 \cdot t'_2)^* \cdot Y(t'_1) \\
&= Y(\widehat{t}) \cdot \left( Y(\widehat{\widehat{t}_1}) \cdot Y(\widehat{t}_2) \cdot Y(t'_2) \right)^* \cdot Y(t'_1) \\
&\sqsubseteq Y(\widehat{t}) \cdot \left( Y(\widehat{\widehat{t}_1}) \cdot Y(\widehat{t}_2)^* \cdot Y(t'_2) \right)^* \cdot Y(t'_1) \\
&= \left\{ \begin{array}{l} Y(\widehat{t}) \cdot Y(\widehat{\widehat{t}_1}) \cdot Y(\widehat{t}_2)^* \cdot Y(t'_2) \cdot Y(t'_1) + \\ Y(\widehat{t}) \cdot \left( Y(\widehat{\widehat{t}_1}) \cdot Y(t'_2) \right)^* \cdot Y(t'_1) \end{array} \right\} \quad (\text{Prop. 5 (2)}) \\
&= \left\{ \begin{array}{l} Y(\widehat{t} \cdot (\widehat{\widehat{t}_1} \cdot \widehat{t}_2^* \cdot t'_2) \cdot t'_1) + \\ Y(\widehat{t} \cdot (\widehat{\widehat{t}_1} \cdot t'_2)^* \cdot t'_1) \end{array} \right\}
\end{aligned}$$

The first expression in this sum equals  $Y((\widehat{t} \cdot \widehat{\widehat{t}_1} \cdot t'_1) \cdot \widehat{t}_2^* \cdot t'_2)$ . This is the pumping yield of the decomposition  $t = (\widehat{t} \cdot \widehat{\widehat{t}_1} \cdot t'_1) \cdot \widehat{t}_2 \cdot t'_2$ . Since  $t_2 = \widehat{t}_2 \cdot t'_2$  is a proper subtree of  $\widehat{t}_1 \cdot t'_1$ , it has fewer nodes than  $\widehat{t}_1 \cdot t'_1$ . So this decomposition is smaller (in the second component), i.e., by induction, the first expression in the above sum is  $\sqsubseteq Y(\mathcal{B}_X)$ .

The second expression in the above sum equals the pumping yield of the decomposition of an  $X$ -tree which is obtained from  $t$  by removing the substructure  $\widehat{t}_2$ . By induction, this pumping yield is  $\sqsubseteq Y(\mathcal{B}_X)$ .  $\square$

## C Proofs of Section 6

**Lemma 1.** For every clean  $\mathbf{g} \in \mathcal{S}[\mathcal{X}]^{\mathcal{X}}$  we can find a system  $\mathbf{f} \in \mathcal{S}[\mathcal{X}']^{\mathcal{X}'}$  ( $\mathcal{X} \subseteq \mathcal{X}'$ ), where  $\mathbf{f}$  is in normal form, and  $\mu \mathbf{g}_X = \mu \mathbf{f}_X$  for all  $X \in \mathcal{X}$ .

*Proof.* For every clean  $\mathbf{g} \in \mathcal{S}[\mathcal{X}]^{\mathcal{X}}$  we can find a  $\mathbf{f} \in \mathcal{S}[\mathcal{X}']^{\mathcal{X}'}$  in normal form with  $\mathcal{X} \subseteq \mathcal{X}'$  such that  $\mu \mathbf{g}_X = \mu \mathbf{f}_X$  for all  $X \in \mathcal{X}$  as follows: We first transform  $\mathbf{g}$  into Chomsky normal-form, which gives us a system  $\mathbf{g}'$  over the same semiring. As the transformation into Chomsky normal-form introduces new variables,  $\mathbf{g}'$  is given in a super set  $\mathcal{X}'$  of  $\mathcal{X}$  with  $\mu \mathbf{g}'_X = \mu \mathbf{g}_X$  for all  $X \in \mathcal{X}$ . Next, as  $\mathbf{g}$  is clean, we can ensure that  $\mathbf{g}'$  is clean, too. We therefore may set  $\mathbf{g}'' := \mathbf{g}' + \mathbf{1}$  without changing the least solution. Hence, every polynomial of  $\mathbf{g}''_X$  has the form

$$c^{(X)} + \sum_{Y, Z \in \mathcal{X}'} a_{Y, Z}^{(X)} \cdot Y \cdot Z \text{ with } 1 \sqsubseteq c \text{ and } a_{Y, Z} \in \{0, 1\}.$$

Finally, as  $\mathbf{1} \sqsubseteq \mu g''$  we have

$$\begin{aligned}
\mu g''_X &= g''_X(\mu g''_X) \\
&= c^{(X)} + \sum_{Y, Z \in \mathcal{X}'} a_{Y, Z}^{(X)} \cdot \mu g''_Y \cdot \mu g''_Z \\
&= c^{(X)} + \sum_{Y, Z \in \mathcal{X}'} a_{Y, Z}^{(X)} \cdot (1 + \mu g''_Y) \cdot (1 + \mu g''_Z) \\
&= c^{(X)} + 1 + \sum_{Y, Z \in \mathcal{X}'} a_{Y, Z}^{(X)} \cdot \mu g''_Y \cdot \mu g''_Z + \sum_{Y \in \mathcal{X}'} \left( \sum_{Z \in \mathcal{X}'} a_{Y, Z}^{(X)} + a_{Z, Y}^{(X)} \right) \cdot \mu g''_Y \\
&= g''_X(\mu g''_X) + 1 + \sum_{Y \in \mathcal{X}'} \left( \sum_{Z \in \mathcal{X}'} a_{Y, Z}^{(X)} + a_{Z, Y}^{(X)} \right) \cdot \mu g''_Y.
\end{aligned}$$

We now define  $f$  by setting for all  $X \in \mathcal{X}'$

$$f_X := g''_X + 1 + \sum_{Y \in \mathcal{X}'} \left( \sum_{Z \in \mathcal{X}'} a_{Y, Z}^{(X)} + a_{Z, Y}^{(X)} \right) \cdot Y.$$


We then have  $g'' \sqsubseteq f$ , and, thus,  $\mu g'' \sqsubseteq \mu f$ , but also  $f(\mu g'') = \mu g''$ , i.e.  $\mu g'' = \mu f$ .  $\square$

For the proof of Theorem 3, we first define partial derivation trees: These trees result from derivation trees by removing exactly one subtree, leaving in some sense a ‘‘dangling pointer’’.

**Definition 8.** Let  $f \in S[\mathcal{X}]^{\mathcal{X}}$ . Let  $t$  be some  $X$ -tree for  $X \in \mathcal{X}$ . Further, let  $Y \in \mathcal{X}$  be some variable such that  $t$  has at least one leaf  $s$  with  $\lambda_v(s) = Y$ . By erasing exactly one such leaf  $s$  from  $t$ , we obtain an  $XY$ -tree. We write  $T_{X, Y}$  for the set of all  $XY$ -trees.

The set  $\mathcal{B}_{X, Y}$  is defined similarly. A tree  $t \in \mathcal{B}_{X, Y}$  results from a tree  $t' \in \mathcal{B}_X$  by removing the exactly one such leaf  $s$  of  $t'$  with  $\lambda_v(s) = Y$ , where  $s$  lies on a longest path from  $t'$  to a leaf.

*Example 3.* Consider the  $X$ -tree depicted on the left. By deleting the leaf labeled by  $(Y, d)$ , we obtain the  $XY$ -tree depicted in the middle, where we represent the missing leaf/subtree by  $\circ$ . Similarly, we obtain the  $XX$ -tree shown on the right by deleting the leaf labeled by  $(X, g)$ .



Note that we can replace  $\circ$  in the  $XY$ -tree by any  $Y$ -tree in order to obtain a valid  $X$ -tree, again. In other words, the yield of an  $XY$ -tree is a linear monomial in  $Y$ .

We now can state our main theorem.

**Theorem 3.** For a finite set of variables  $\mathcal{X}$  ( $n := |\mathcal{X}|$ ), let  $\mathcal{S}$  be a lossy semiring, and  $\mathbf{f} \in \mathcal{S}[\mathcal{X}]^{\mathcal{X}}$  a clean and strongly connected system of polynomials in normal-form. We then have  $\mu\mathbf{f} = \mu\mathbf{f}_{\mathcal{B}}$ .

*Proof.* We again show that we can transform any  $X$ -tree  $t$  w.r.t.  $\mathbf{f}$  into a tree  $\hat{t}$  contained in  $\mathcal{B}_X$  with  $Y(t) \sqsubseteq Y(\hat{t})$ . We proceed by induction on the number  $N$  of nodes of  $t$ . If  $N = 1$ , then  $t$  has height 0. By definition, we have  $t \in \mathcal{B}_X$ , so we are done.

Therefore assume  $N > 1$ . As  $\mathbf{f}$  is in normal form, we either have  $\lambda_m(t) = b_l Y b_r$  or  $\lambda_m(t) = YZ$  for some  $Y, Z \in \mathcal{X}$ , and  $b_l, b_r \in \mathcal{S} \setminus \{0\}$ . If  $t$  is labeled by  $\lambda_m(t) = b_l Y b_r$ ,

$$\begin{array}{c} t = (X, b_l Y b_r) \\ | \\ t_1 \end{array}$$

then  $t$  has exactly one child  $t_1$ , which immediately can be replaced by some tree  $\hat{t}_1$  in  $\mathcal{B}_Y$  with  $Y(t_1) = Y(\hat{t}_1)$  because of induction. This gives us the tree  $\hat{t}$

$$\begin{array}{c} \hat{t} = (X, b_l Y b_r) \\ | \\ \hat{t}_1 \end{array}$$

and  $Y(\hat{t}) = b_l Y(\hat{t}_1) b_r = b_l Y(t_1) b_r = Y(t)$ .


Hence, assume that  $\lambda_m(t) = YZ$ , i.e.  $t$  has two children  $t_1, t_2$ .

$$\begin{array}{c} t = (X, YZ) \\ / \quad \backslash \\ t_1 \quad t_2 \end{array}$$


Descending into  $t_1$  by always taking the left most child, we end up at the left most leaf  $s$  of  $t$ . We denote by  $t_{1,1}$  to  $t_{1,k}$  the “right” children of the nodes located on the path from  $t_1$  to  $s$  for some  $k \in \mathbb{N}$ . Let  $r$  then be the father of  $s$  with  $\lambda_v(s) = V$ , and  $\lambda_m(s) = a \in \mathcal{S}$ . We assume that  $\lambda_m(r) = VW$  for some  $W \in \mathcal{X}$

$$\begin{array}{c} t = (X, YZ) \\ / \quad \backslash \\ t_1 \quad t_2 \\ / \quad \backslash \\ r = (U, VW) \quad t_{1,1} \\ / \quad \backslash \\ s = (V, a) \quad t_{1,k} \end{array}$$


As  $\mathbf{f}$  is in normal form, and  $VW$  is a monomial of  $\mathbf{f}_U$ , there exists also a monomial  $c_l W c_r$  appearing in  $\mathbf{f}_U$  for some  $c_l, c_r \in \mathcal{S} \setminus \{0\}$ . We first remove from  $t_1$  the leaf  $s$ , and relabel the node  $r$  by setting  $\lambda_m(r) := c_l W c_r$ . This gives us the tree  $t'_1$  with  $Y(t_1) \sqsubseteq a \cdot Y(t'_1)$ , as  $1 \sqsubseteq c_l, c_r$ :




Now, as  $f$  is strongly connected and in normal form, we find an  $Y$ -tree  $t_Y$  of height at most  $n - 1$  which has  $(V, a)$  as its single leaf, such that  $a \sqsubseteq Y(t_Y)$ ; similarly, we find a  $ZX$ -tree  $t_{ZX}$  of height at most  $n - 1$  having  $\circ$  as its single leaf; the “yield” of  $t_{ZX}$  is some monomial  $d_l X d_r$  for some  $d_l, d_r \in S \setminus \{0\}$ . Using these, we construct the following tree  $t'$  with  $\lambda_v(t') = X$ , and  $\lambda_m(t') = YZ$ . As left child of  $t'$ , we take the  $Y$ -tree  $t_Y$ , whereas we take  $t_{ZX}$  as the right child, giving us:



We next replace  $\circ$  by a new node  $t''$  with  $\lambda_v(t'') = X$ , and  $\lambda_m(t'') = YZ$ :




Finally, we set the  $Y$ -tree  $t'_1$  as the left child of  $t''$ , while the  $Z$ -tree  $t_2$  becomes the right child of  $t''$ :




We now have  $Y(t') = Y(t_Y) \cdot Y(t_{ZX}) \sqsupseteq a \cdot d_l \cdot Y(t'') \cdot d_r \sqsupseteq a \cdot Y(t'_1) \cdot Y(t_2) \sqsupseteq Y(t_1) \cdot Y(t_2) = Y(t)$ .

By construction of  $t'$ , the left child is a  $Y$ -tree of height at most  $n - 1$ , while every node from  $t_{ZX}$  to  $t''$  has exactly one child. Hence, only the subtree  $t''$  might not have the required form. But as  $t''$  has one node less than  $t$ , we find by induction on the number of nodes a tree  $\hat{t}'' \in \mathcal{B}_X$  with  $Y(\hat{t}'') \sqsubseteq Y(t'')$ . Replacing in  $t'$  the subtree  $t''$  by this tree  $\hat{t}''$ ,




we then obtain the tree  $\hat{t}$  with  $\hat{t} \in \mathcal{B}_X$  and  $Y(\hat{t}) \sqsupseteq Y(t') \sqsupseteq Y(t)$ .


If  $\lambda_m(r) = c_l V c_r$  for some  $c_l, c_r \in S \setminus \{0\}$ , i.e.



we proceed similarly, but we define  $t'_1$  as follows: again, we remove the leaf  $s$  from  $t_1$ , but as  $r$  has  $s$  as its only child, we now relabel  $r$  by  $\lambda_m(r) := \mathbf{f}_U(\mathbf{0})$ . As  $\mathbf{f}$  is clean, we have  $\mathbf{f}_U(\mathbf{0}) \sqsupseteq 1$ . This gives us:



and



Again, we can find a  $\hat{t}'' \in \mathcal{B}_X$  with  $Y(\hat{t}'') \sqsubseteq Y(t'')$  as  $t''$  has one node less than  $t$ , and the induction is complete.  $\square$

## D Proofs of Section 7

**Theorem 5.**  $\mu f = f^n(\mathbf{0})$  holds for polynomial systems over 1-bounded semirings.

*Proof.* We reuse the notation from the proof of Theorem 2: If  $t_2$  is a subtree of a derivation tree  $t$ , we write  $t = t_1 \cdot t_2$  where  $t_1$  is the partial derivation tree obtained from  $t$  by removing  $t_2$ .

Recall that, by Proposition 2,  $(f^n(\mathbf{0}))_X = Y(\mathcal{T}_X^{(n-1)})$ , where  $\mathcal{T}_X^{(n-1)}$  contains all  $X$ -trees of height at most  $n - 1$ . We proceed by derivation tree analysis, i.e., by discharging the precondition of Proposition 3. So it suffices to show that for any  $X$ -tree  $t$  there is a tree  $t'$  of height at most  $n - 1$  with  $Y(t) \sqsubseteq Y(t')$ . We proceed by induction on the number of nodes in  $t$ . For the induction base,  $t$  has just one node, so  $t \in \mathcal{T}_X^{(0)}$ . For the induction step w.l.o.g. let  $t$  be an  $X$ -tree with a height of at least  $n$ . Then there is a decomposition  $t = t_1 \cdot t_2 \cdot t_3$  with  $\lambda_1(t_2) = \lambda_1(t_3)$ . We have  $Y(t) = y_1 y_2 y_3 y_4 y_5$  where  $Y(t_1) = y_1 y_5$ ,  $Y(t_2) = y_2 y_4$  and  $Y(t_3) = y_3$ . Let  $t' = t_1 \cdot t_3$ . Notice that  $t'$  is a valid  $X$ -tree as  $\lambda_1(t_2) = \lambda_1(t_3)$ . We have  $Y(t') = y_1 y_3 y_5$  which is, by 1-boundedness, at least  $y_1 y_2 y_3 y_4 y_5 = Y(t)$ . As  $t'$  has fewer nodes than  $t$ , there is, by induction hypothesis, an  $X$ -tree  $t''$  of height at most  $n - 1$  such that  $Y(t') \sqsubseteq Y(t'')$ . Combined we get  $Y(t) \sqsubseteq Y(t') \sqsubseteq Y(t'')$ .  $\square$